White, M., Bush, J., Kai, J., Bhopal, R., & Rankin, J. (2006). Quitting smoking and experience of smoking cessation interventions among UK Bangladeshi and Pakistani adults: The views of community members and health professionals. *Journal of Epidemiology and Community Health*, 60, 405–411.

Acknowledgment. This study was supported with financial support to the first author by the Foundation of the University of Medicine and Dentistry of New Jersey (grant # 28–06) and the University of Medicine and Dentistry of New Jersey–School of Nursing.

Correspondence regarding this article should be directed to Sun S. Kim, PhD, RN, University of Massachusetts Medical School, Department of Psychiatry, 55 Lake Avenue North, Suite 57-412, Worcester, MA 01655. E-mail: sun.kim@umassmed.edu

APPENDIX

Comparison of the Minnesota Tobacco Withdrawal Scale Between the Original and a Korean Version

The Original Version						A Korean Version					
Minnesota Tobacco Withdrawal Scale Please rate your experience of the following nine symptoms.						미네소타 담배 금단증상 측정기 다음 9 개의 증상들 중 귀하의 경험을 가장 잘 나타내고 있는 곳에 동그라미 표하세요.					
Not Present = NP; Slight = SL; Mild = MI; Moderate = MO; Severe = SE						전혀 없다 = 전혀; 아주 조금 = 아조; 약하게 = 약; 보통으로 = 보통; 심하게 = 심					
1. Craving for cigarettes or desire to smoke	NP	SL 1	MI	МО	SE	1. 담배를 피고 싶은 절실함	전혀	아조	약	보통	심
2. Irritability/frustration /anger	NP	SL	MI	MO	SE	2. 짜증남/좌절/분노	진혀	아조	약	보통	심
3. Anxiety	NP	SL	MI	MO	SE	3. 불안	전혀	아조	약	보통	심
4. Difficulty concentrating	NP	SL	MI	MO	SE	4. 집중하기 어려움	전혀	아조	약	보통	심
5. Restlessness	NP	SL	MI	MO	SE	5. 안절부절 못함	전혀	아조	약	보통	심
6. Increased appetite	NP	SL	MI	MO	SE	6. 식욕 증가	전혀	아조	약	보통	심
7. Disturbed sleep	NP	SL	MI	MO	SE	7. 수면 방해	전혀	아조	약	보통	심
8. Depression	NP	SL	MI	MO	SE	8. 우울함	전혀	아조	약		심
9. Impatience	NP	SL	MI	MO	SE	9. 참지 못함	전혀	아조	약	보통	심